

PCTWORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁵: A61K 31/01, 31/02, 31/015, 31/19, 31/045, 31/265	A1	(11) International Publication Number: WO 94/28883 (43) International Publication Date: 22 December 1994 (22.12.94)
(21) International Application Number: PCT/US94/06653 (22) International Filing Date: 8 June 1994 (08.06.94) (30) Priority Data: 073,934 8 June 1993 (08.06.93) US (60) Parent Application or Grant (63) Related by Continuation US 073,934 (CON) Filed on 8 June 1993 (08.06.93) (71)(72) Applicant and Inventor: BROWN, Raymond, K. [US/US]; 140 West 69th Street, New York, NY 10023 (US). (74) Agents: ROBINSON, Joseph, R. et al.; Darby & Darby, 805 3rd Avenue, New York, NY 10022 (US).	(81) Designated States: AT, AU, BB, BG, BY, CA, CH, CN, CZ, DE, DK, ES, FI, GB, GE, HU, JP, KG, KP, KR, KZ, LK, LU, LV, MD, MG, MN, MW, NL, NO, NZ, PL, RO, RU, SD, SE, SI, SK, TJ, TT, UA, US, UZ, VN. Published <i>With international search report.</i>	
(54) Title: THERAPEUTIC COMPOSITIONS AND METHODS OF USE		
(57) Abstract According to the present invention, there is provided an admixture of sorbic acid, malic acid, fumaric acid, crotonic acid, and optionally aconitic acid. Further contemplated by the present invention are methods for treating various pathological conditions such as viral infections, acidosis, tumors, and bacterial and fungal infections by administering various therapeutically effective amounts of an admixture described above.		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	GB	United Kingdom	MR	Mauritania
AU	Australia	GE	Georgia	MW	Malawi
BB	Barbados	GN	Guinea	NE	Niger
BE	Belgium	GR	Greece	NL	Netherlands
BF	Burkina Faso	HU	Hungary	NO	Norway
BG	Bulgaria	IE	Ireland	NZ	New Zealand
BJ	Benin	IT	Italy	PL	Poland
BR	Brazil	JP	Japan	PT	Portugal
BY	Belarus	KE	Kenya	RO	Romania
CA	Canada	KG	Kyrgyzstan	RU	Russian Federation
CF	Central African Republic	KP	Democratic People's Republic of Korea	SD	Sudan
CG	Congo	KR	Republic of Korea	SE	Sweden
CH	Switzerland	KZ	Kazakhstan	SI	Slovenia
CI	Côte d'Ivoire	LI	Liechtenstein	SK	Slovakia
CM	Cameroon	LK	Sri Lanka	SN	Senegal
CN	China	LU	Luxembourg	TD	Chad
CS	Czechoslovakia	LV	Latvia	TG	Togo
CZ	Czech Republic	MC	Monaco	TJ	Tajikistan
DE	Germany	MD	Republic of Moldova	TT	Trinidad and Tobago
DK	Denmark	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	US	United States of America
FI	Finland	MN	Mongolia	UZ	Uzbekistan
FR	France			VN	Viet Nam
GA	Gabon				

5

THERAPEUTIC COMPOSITIONS AND METHODS OF USE10 Field of the Invention

The invention relates to admixture compositions of short chain organic fatty acids which have biological effects. For example, they are useful in the treatment of numerous pathologies that affect mammals. The invention also relates to methods of treatment of such pathologies, wherein appropriate therapeutically effective amounts of the compositions are administered.

Background of the Invention

A wide variety of carboxylic acid containing compositions are biologically active, and several short chain organic acids which are naturally occurring acids have previously been implicated in the treatment of various pathological conditions. For example, Nordman, U.S. Patent No. 3,291,689, discloses the treatment of hepatic ammonia intoxication with a mixture of L-arginine and malic acid. U.S. Patent No. 3,718,664 discloses the use of thioctic acid derivatives in the treatment of acidosis.

Sloan, U.S. Patent No. 4,381,307, discloses soft tertiary amine ester derivatives that have biological effects, while U.S. Patent No. 4,760,078 discloses a 1,2, dithiol-3-thione derivative that has an immunomodulating effect.

Rubenstein, U.S. Patent No. 4,971,760, discloses the sterilization of blood, blood constituents, and transplant tissues with a disinfectant that includes lactic acid and sodium chlorite. Naphtholic acid derivatives were discovered to be useful in enhancing oxygen availability to mammalian tissue by Suh et al., U.S. Patent No. 5,015,663.

Hoffler et al., U.S. Patent No. 5,043,357, disclose a virucidal agent that is predominantly comprised of ethanol and/or alcohol but which includes a minor amount of a short chain organic acid. Further, U.S. Patent No. 5,093,140 discloses an aqueous bactericide that contains organic acids. This solution is used in the scalding or washing stages of meat dressing. Kross et al., U.S. Patent No. 5,100,652 disclose an oral disinfectant that contains organic acids. See also Comroe et al., *The Lung*, 1955 Yearbook of Medical Publications Inc., Chptr. 4; John West, *Respiratory Physiology*, The Williams & Wilkins Co., Chptr. 6; *Hypoxia, Metabolic Acidosis and Circulation*, ARTCF...Oxford; Arnold et al, "Excessive Intracellular Acidosis of Skeletal Muscle on Exercise in a Patient with a Post-Viral Exhaustion/Fatigue Syndrome" *The Lancet*: 6/23/84; Schweckendiek, W., "Heilung von Psoriasis vulgaris", *Med. Sschr*, 13:103-4, 1959; G.E. Abraham et al., "Rationale for the Use of Magnesium and Malic Acid", *Journal of Nutritional Medicine*, 3:49-49 (1992); Kuroda, Z. and Akano, M., "Antitumor and Anti-intoxication Activities of Fumaric Acid in Cultured Cells", *Gann*, 727:77-782 (1981); *Chemical Abstracts* 98945Y, 116, Mar 1992 "Antifungal Activity of Fumarates in Mice Infected with *C. Albicans*"; Bauer et al., "Clinische Wochenschrift 1991 69(2), pp. 722-4; for further details of several pathological processes which are susceptible to treatment according to the present invention.

It has now been discovered that admixtures of specific short chain organic acids have broad therapeutic effects.

Summary of the Invention

According to the present invention, there is provided a composition which comprises an admixture of sorbic acid, aconitic acid, malic acid, fumaric acid, crotonic acid, and optionally aconitic acid.

Further contemplated by the present invention are methods for treating various pathological conditions such as viral infections including, but not limited to, cytomegalovirus and hepatitis infections; acidosis; tumors; and bacterial and

fungal infections, by administering various therapeutically effective amounts of the admixture described above.

Detailed Description of the Invention

Organic acids are a large group of compounds, many of which can be derived from proteins, carbohydrates, and fats. Although many occur naturally, many can be synthetically produced without great difficulty.

Sorbic acid is a crystalline diolefinic acid having the formula $C_6H_8O_2$. Typically, it is used as mold and yeast inhibitor; a fungistatic agent for foods, especially cheeses; or to improve the characteristics of drying oils, the gloss of alkyd type coatings, or the milling properties of cold rubber.

Aconitic acid is a white crystalline acid having the formula $C_6H_6O_6$. Typically, it is used in manufacturing itaconic acid or as a plasticizer for buna rubber and plastics.

Malic acid is a crystalline dicarboxylic acid having the formula $C_4H_4O_5$, and is typically found in three isomeric crystalline forms.

Fumaric acid is a crystalline unsaturated dicarboxylic acid having the formula $C_4H_4O_4$. Fumaric acid is typically used as a substitute for tartaric acid in beverages and baking powders, as a replacement or partial replacement of citric acid in soft drinks, and as an antioxidant.

Crotonic acid is an unsaturated aliphatic acid having the formula $C_4H_6O_2$. It exists in the crystalline form, and is typically used in the manufacture of copolymers with vinylacetate that are used in lacquers and paper sizing, and in the manufacture of softening agents for synthetic rubber.

It has now been discovered that these acids, in combination, have significant therapeutic value. The admixtures of these acids are prepared by conventional means known to those skilled in the skilled art. The order of addition of the acids to the admixture does not affect the final composition.

Although a wide range of proportions of the individual components of the admixture is effective, in a typical

composition, each of the four required individual acids will independently comprise from about 1 to about 50 parts by weight and, optionally, aconitic acid will comprise from 0 to about 50 parts by weight, based upon 100 parts by weight of the five acids combined. Preferably, sorbic acid comprises from about 5 to about 40 parts by weight, aconitic acid comprises from 0 to about 50 parts by weight, malic acid comprises from about 1 to about 30 parts by weight, fumaric acid comprises from about 1 to about 30 parts by weight, and crotonic acid comprises from about 1 to about 30 parts by weight, based upon 100 parts by weight of the five acids combined. Most preferably, sorbic acid comprises from about 20 to about 30 parts by weight, aconitic acid comprises from about 35 to about 45 parts by weight, malic acid comprises from about 10 to about 20 parts by weight, fumaric acid comprises from about 10 to about 20 parts by weight, and crotonic acid comprises from about 1 to about 20 parts by weight, based upon 100 parts by weight of the five acids combined. Special mention is made of an admixture wherein sorbic acid comprises about 24 parts by weight, aconitic acid comprises about 40 parts by weight, malic acid comprises about 16 parts by weight, fumaric acid comprises about 16 parts by weight, and crotonic acid comprises about 4 parts by weight, based upon 100 parts by weight of the five acids combined.

Suitable pharmaceutically active carriers may be combined with the admixture. Such carriers include those known to those skilled in the art and can be added at any point in the mixing process. If a pharmaceutically active carrier is used, the carrier will comprise from about 1 to about 99 parts by weight and the acid admixture will comprise from about 99 to about 1 part by weight, based upon 100 parts by weight of acid admixture and carrier combined.

The acid admixture with or without the carrier can be formulated into dosage unit forms including, but not limited to, capsules and tablets. Methods of preparation of dosage unit forms would be known to those skilled in the art. Agents which

facilitate the manufacture and/or use of such dosage unit forms may be added such as plasticizers, lubricants, excipients, diluents and the like.

Methods of treatment of various pathologies contemplated by the present invention involve the administration of therapeutically effective amounts of the admixture to mammals in need of such treatment. Administration is by means known to those skilled in the art. Preferably, administration is systemic and most preferably, it is oral.

Pathologies which are effectively treated with the organic acid admixture described herein include but are not limited to acidosis; viral infections including, but not limited to, cytomegalovirus and hepatitis infections; tumors; and bacterial and fungal infections, including but not limited to S. aureus, E. coli, and C. albicans.

All of these pathologies are well known to those skilled in the art, and their diagnosis would be evident to one skilled in the art. For example, acidosis is reflected by abnormally acidic blood pH and/or urine pH. Normal blood pH ranges from about 7.3 to about 7.38, and normal urine pH ranges from about 6 to about 8. Viral, bacterial, and fungal infections can be confirmed by appropriate assay methods. Other diseases such as chronic fatigue syndrome are only diagnosed through clinical observations. Various opportunistic infections which are characteristic of AIDS including, but not limited to, cytomegalovirus, hepatitis, and the like are effectively treated as described herein.

The amount of the admixture necessary to treat acidosis is an anti-acidosis effective amount. Similarly, the amounts of the admixture necessary to treat a viral infection, and particularly hepatitis and cytomegalovirus infections are an anti-viral effective amount, and particularly anti-hepatitis and an anti-cytomegalovirus effective amounts. The amount required to treat tumors, and bacterial and fungal infections are anti-

tumor, anti-bacterial, and anti-fungal effective amounts, respectively.

The actual amounts of the admixture to be administered will depend independently upon the age, weight, sex, sensitivity, medical condition, including but not limited to stage of a particular infection or disease, or the like, of an individual. However, the amount will be a safe non-toxic amount.

These amounts can be determined by experimentation well-known in the art such as by establishing a matrix of dosages and frequencies and assigning a group of experimental subjects to each point in the matrix. Typically, that amount will range from about 1000 to about 2000 milligrams per day for a 150 pound man and preferably about 1500 milligrams per day for a 150 pound man. Appropriate dosages for different body weights can be calculated from this.

A daily dosage identifies the average amount of the mixture administered to an individual. Although the daily dosage may actually be administered daily, it need not be administered daily. The daily dosage is merely an average dosage that an individual receives when the mixture is administered over a period of time. The daily dosage can be administered in divided portions so that the total amount administered is the daily dosage.

Description of the Preferred Embodiments

The following examples illustrate the invention without limitation. All amounts are given as parts by weight (pbw).

Example 1

An admixture of 24 parts by weight of sorbic acid, 40 parts by weight of aconitic acid, 16 parts by weight of malic acid, 16 parts by weight of fumaric acid, and 4 parts by weight of crotonic acid was applied, in various dilutions, to cut filter paper, and the filter paper was overlaid on plates streaked with bacteria. Results are illustrated in Table 1.

Table 1 - Bacterial Assays						
<u>Admixt</u> <u>ure</u> <u>Diluti</u> <u>on</u>	<u>Aspergi</u> <u>llus</u>	<u>Staph.</u> <u>aureus</u>	<u>Pseudom</u> <u>onas</u> <u>aerugin</u> <u>osa</u>	<u>Escheri</u> <u>chia</u> <u>coli</u>	<u>Candida</u> <u>albican</u> <u>s</u>	<u>pH</u>
Full Streng th	-	1.6 cm	-	1.1 cm	1.6 cm	2.2
1:10	-	-	-	-	-	4.6
1:100	-	-	-	-	-	5.2

The (-) negative means that no inhibition was observed.

Example 2

A patient had a temperature of 101°F and symptoms of acute urinary and prostate infection. The patient was treated with 1500 mg/day of an admixture of 24 pbw of sorbic acid, 40 pbw of aconitic acid, 16 pbw of malic acid, 16 parts by weight of fumaric acid, and 4 pbw of crotonic acid based upon 100 pbw of total admixture for ten days. Symptoms and temperature subsided by the 4th or 5th day. Antimicrobial and alkalizing activity and therapeutic effects of the admixture are illustrated in Table 2. Cultures which were positive for streptococci were negative by the third week.

Table 2 - Urinary Infection			
Laboratory Test	Initial Treatment Day 1	Day 10	Day 32
5 URINALYSIS			
pH	5.5	8	8
Protein	1+	0	0
WBC	20-30	2-4	2-4
Bacteria	many	0	few
10 Urine Culture	Strept. +	±	0
Blood PSA (normal 0-4)	21.9	5.1	3.22
WBC	10.4		6.1

15 Example 3

Six HIV positive individuals with average urine pH at breakfast, lunch, and dinner of 6.0, as measured by pHDrion tape, were administered 1500 mg/day of an admixture of 24 pbw of sorbic acid, 40 pbw of aconitic acid, 16 pbw of malic acid, 16 pbw of fumaric acid, and 4 pbw of crotonic acid.

After one month, average urine pH was raised to 6.3. Previously, it was fixed at 6.0. Results are illustrated in Table 3 below.

Table 3 - Acidosis			
Patient	Urine pH Breakfast	Urine pH Lunch	Urine pH Dinner
"A"	6.0	6.2	6.2
	6.0	6.2	6.0
	6.4	6.2	6.2
"B"	6.2	6.4	6.2
	6.0	6.0	6.4
	6.2	6.4	6.2
"C"	6.0	6.0	6.2
	6.0	6.2	6.0
	6.0	6.0	6.2
"D"	6.0	6.0	6.2
	6.0	6.2	6.2
	6.0	6.0	6.2
"E"	6.0	6.4	6.4
	6.2	6.2	6.2
	6.0	6.0	6.2
"F"	6.0	6.4	6.4
	6.2	6.2	6.2
	6.0	6.2	6.2

Example 4

A patient with urine culture positive for cytomegalovirus infection was treated with 1500 mg/day of an admixture of 24 pbw of sorbic acid, 40 pbw of aconitic acid, 16 pbw of malic acid, 16 pbw of fumaric acid, and 4 pbw of crotonic acid, based upon 100 pbw of total admixture for six weeks. After six weeks, a cytomegalovirus urine culture was negative.

Example 5

A patient having cancer of the prostate with a PSA of 17 was treated with Zolodex™. PSA was lowered to 2.4. Seven months after Zolodex treatment was discontinued, PSA rose to 13.

The patient then received 1500 mg/day of an admixture of 24 pbw of sorbic acid, 40 pbw of aconitic acid, 16 pbw of malic acid, 16 pbw of fumaric acid, and 4 pbw of crotonic acid based upon 100 pbw of total admixture, and PSA was lowered to

1.65. Nightly urination decreased from 5-6 times/night to two times/night and the patient's general energy increased 80%.

Example 6

5 A patient was diagnosed with a pancreatic mass. The patient also had nausea, pain, an inability to eat, weight loss, and weakness.

10 The patient, who weighed about less than eighty pounds, was given 50 mg/day of an admixture of 24 pbw of sorbic acid, 40 pbw of aconitic acid, 16 pbw of malic acid, 16 pbw of fumaric acid, and 4 pbw of crotonic acid based upon 100 pbw of total admixture for two months. The patient's nausea abated, her appetite improved, she slept more restfully, and she did not urinate as frequently at night. The epigastric mass was no longer palpable.

15 Example 7

20 A patient underwent a resection of a recurrent bladder tumor, at which time, the pathological specimen revealed Grade II/III moderately differentiated papillary transitional cell carcinoma with invasion of the muscularis of the urinary bladder and of the prostate. The patient was placed on a protocol of 1500 mg/day of an admixture of 24 pbw of sorbic acid, 40 pbw of aconitic acid, 16 pbw of malic acid, 16 pbw of fumaric acid, and 4 pbw of crotonic acid based upon 100 pbw of total admixture. A repeat cystoscopy two months later showed only several small
25 papillary lesions in the urinary bladder and a small lesion in the prostatic urethra. Rectal examination was unremarkable. CAT SCAN confirmed the improvement observed by cystoscopy.

Example 8

30 A patient was diagnosed with long standing Hepatitis C. The patient was treated with 1500 mg/day of an admixture of 24 pbw of sorbic acid, 40 pbw of aconitic acid, 16 pbw of malic acid, 16 pbw of fumaric acid, and 4 pbw of crotonic acid based upon 100 pbw of total admixture. After one month, pain in the patient's side subsided. The patient was no longer nauseated and

generally felt better. His SGOT laboratory value went down from 68 to 37.

Example 9

An admixture of 24 pbw of sorbic acid, 40 pbw of aconitic acid, 16 pbw of malic acid, 16 pbw of fumaric acid, and 4 pbw of crotonic acid based upon 100 pbw of total admixture was tested for virucidal effectiveness against Cytomegalovirus. The viral inoculum was diluted 1:10 with the admixture. The test material/virus mixture was sampled at 3 and 6 minutes. The treated viral mixture was neutralized and titered. An initial viral titer was determined.

The challenge organism was Cytomegalovirus, (CMV) and the host cell line was MRC-5, ATCC CCL-171. Reagents and media were EMEM with Earle's salts supplemented with heat inactivated fetal bovine serum (FBS), glutamine, and penicillin streptomycin (MEM complete); sterile phosphate buffered saline; and newborn calf serum (NBCS).

The virus stock culture was titered and adjusted to contain approximately 10^7 CCID₅₀/ml (50% cell-culture infectious doses per milliliter) and stored in liquid nitrogen until use. The virus stock was thawed and was diluted 10-fold in MEM through 10^7 . One ml aliquots of appropriate dilutions were plated onto 5 confluent MRC-5 monolayers. The cells were incubated at $37 \pm 1^\circ\text{C}$, in 5% CO₂, for 2 hours to allow the virus to adsorb to the cells. After adsorption, the medium was withdrawn from the monolayers, the cells were washed once with PBS, and refed with MEM. The monolayers were incubated for 14 days at $37 \pm 1^\circ\text{C}$, in 5% CO₂, and subsequently were observed for virus-specific cytopathogenic effects (CPE).

The test solution was tested by combining the virus with the admixture at a 1:10 dilution. This suspension was agitated continuously at a constant temperature of 25°C and samples were withdrawn at 3 and 6 minutes to determine a survivor curve and to calculate a D-value.

The sampled suspensions were neutralized with a 1:1 dilution in new-born calf serum and were diluted ten-fold in MEM. One ml aliquots of the neutralized mixture were plated onto MRC-5 monolayers (four wells per dilution) and were incubated at $37 \pm 1^\circ\text{C}$ in a 5% CO_2 in air atmosphere for 2 hours. Following the incubation period, the fluids were removed. The monolayers were washed with PBS, refed MEM and were incubated at $37 \pm 1^\circ\text{C}$ in a 5% CO_2 in air atmosphere for thirteen days.

Control virus stock was titered at the time of the test as described, to confirm that the titer was in an acceptable range. The maintenance of the virus titer was evaluated using PBS instead of the test solution. Samples were taken less than 15 seconds after the initiation of the test (zero time control) and after 6 minutes (final time control). The average $\text{CCID}_{50}/\text{ml}$ for each titer was determined by the Reed and Muench method. *American Journal of Hygiene*, 1938, 27:493. The D-value was calculated by the Stumbo method. Stumbo, C.R., *Thermobacteriology in food processing*, New York Academic Press (1973), pp. 235-247

No CPE was observed at 3 minutes. A D-value calculated using this time resulted in a D-value of 45 seconds. This is an estimate since the first sampling was done at 3 minutes (no data were collected at less than 3 minutes).

When a survivor curve was determined in this test, the admixture provided data used to calculate a D-value of 45 seconds (estimated). A conclusion can be drawn that the admixture was antiviral against cytomegalovirus.

Results are illustrated in Tables 4, 5, and 6 below.

Table 4					
Test and Control Titers Expressed as CPE Day 13 Results					
		PBS		TEST SOLUTION	
DILUTION	VIRUS STOCK	Zero	6 min	3 min	6 min
10 ⁻¹	NA	++++	++++	----	----
10 ⁻²	++++	++++	++++	----	----
10 ⁻³	++++	++++	++++	----	----
10 ⁻⁴	++++	+++-	+++-	----	----
10 ⁻⁵	++++	----	----	----	----
10 ⁻⁶	----	----	----	----	----
10 ⁻⁷	----	N/A	N/A	N/A	N/A
CCID ₅₀ /ml	3.2 x 10 ⁵	1 x 10 ⁴	1 x 10 ⁴		

+ = CPE observed
 - = No CPE observed
 CT = Cytotoxicity observed
 0 = No Cytotoxicity observed

Table 5	
Control and Test Titers at Time Intervals After Exposure to Scam Disinfectant	
<u>SAMPLE</u>	<u>CCID₅₀/ml</u>
Sample	3.2 x 10 ⁵
PBS at Zero time	1 x 10 ⁴
PBS at 6 minutes	1 x 10 ⁴
3 minutes	No CPE observed
6 minutes	No CPE observed

Table 6	
Determination of the D-Value	
5	$D = \frac{T_x - T_o}{\log A_o - \log A_x} = \frac{3 - 0}{\log 10^4 - 0} = \frac{3}{4} = 0.75 \text{ min} = 45 \text{ seconds}$

- Tx = end of time interval
 To = beginning of time interval
 10 AO = CCID₅₀/ml at To
 AX = CCID₅₀/ml at Tx

Example 10

A 36 year old male with well controlled HIV infection
 15 had progressively deteriorating kidney function from no
 demonstrable cause. He averaged 5x nocturia and ran a
 consistently 3 plus albuminuria with urinary specific gravity
 fixed at 1.005 (normal to 1.030).

After approximately 4 months of 1500 mg/day of
 20 admixture of 24 pbw of sorbic acid, 40 pbw of aconitic acid, 16
 pbw of malic acid, 16 pbw of fumaric acid, and 4 pbw of crotonic
 acid based upon 100 pbw of total admixture, his kidney function
 normalized. His three plus albuminuria gradually became
 negative; he became able to sleep through nights, and the
 25 specific gravity of his urine by concentration test increased to
 1.0020. His urinary pH, which had been fixed at 5-6 resumed a
 normal pattern of morning aciduria, increasing to neutral or
 alkaline during the day according to his diet and activity.

Accompanying this was a total regression of his long-
 30 existing bilateral massively swollen parotid glands; the non-
 specific diagnosis derived from multiple biopsies had been cystic
 parotitis. The presumptive retro-diagnosis for parotitis and the
 kidney insufficiency was severe cytomegalovirus (CMV) infection,
 suggested by high titers of CMV antibodies.

35 The above mentioned patents, test methods, and
 publications are hereby incorporated by reference in their
 entirety.

Many variations of the present invention will suggest themselves to those skilled in the art in light of the above detailed description. All such obvious variations are within the full intended scope of the appended claims.

IN THE CLAIMS

1. A composition comprising an admixture comprising:
 - (a) sorbic acid;
 - (b) malic acid;
 - (c) fumaric acid;
 - (d) crotonic acid; and
 - (e) optionally, aconitic acid.
2. A composition as defined in claim 1 further comprising (f) a pharmaceutically acceptable carrier.
3. A composition as defined in claim 1, wherein each of components (a), (b), (c), and (d) independently comprise from about 1 to about 50 parts by weight, and component (e) comprises from 0 to about 50 parts by weight, based upon 100 parts of by weight of (a), (b), (c), (d), and (e) combined.
4. A composition as defined in claim 3, wherein component (a) comprises from about 5 to about 40 parts by weight, component (b) comprises from about 1 to about 30 parts by weight, component (c) comprises from about 1 to about 30 parts by weight, and component (d) comprises from about 1 to about 30 parts by weight, and component (e) comprises from 0 to about 50 parts by weight, based upon 100 parts by weight of components (a), (b), (c), (d) and (e) combined.
5. A composition as defined in claim 4, wherein component (a) comprises from about 20 to about 30 parts by weight, component (b) comprises from about 10 to about 20 parts by weight, component (c) comprises from about 10 to about 20 parts by weight, component (d) comprises from about 1 to about 20 parts by weight, and component (e) comprises from about 35 to about 45 parts by weight, based upon 100 parts by weight of components (a), (b), (c), (d) and (e) combined.

1 6. A composition as defined in claim 5, wherein
2 component (a) comprises about 24 parts by weight, component (b)
3 comprises about 16 parts by weight, component (c) comprises about
4 16 parts by weight, component (d) comprises about 4 parts by
5 weight, and component (e) comprises about 40 parts by weight,
6 based upon 100 parts by weight of components (a), (b), (c), (d),
7 and (e) combined.

1 7. A method for treating acidosis in a mammal in need
2 of such treatment comprising administering to said mammal an
3 anti-acidosis effective amount of a composition as defined in
4 claim 1.

1 8. A method for treating a viral infection in a
2 mammal in need of such treatment comprising administering to said
3 mammal an anti-viral effective amount of a composition as defined
4 in claim 1.

1 9. A method for treating cytomegalovirus infection in
2 a mammal in need of treatment comprising administering to said
3 mammal an anti-cytomegalovirus effective amount of a composition
4 as defined in claim 1.

1 10. A method for treating hepatitis in a mammal in
2 need of such treatment comprising administering to said mammal an
3 anti-hepatitis effective amount of a composition as defined in
4 claim 1.

1 11. A method for treating a tumor in a mammal in need
2 of such treatment comprising administering to said mammal an
3 anti-tumor effective active amount of a composition as defined in
4 claim 1.

1 12. A method for treating a bacterial infection in a
2 mammal in need of such treatment comprising administering to said

3 mammal an anti-bacterial effective amount of a composition as
4 defined in claim 1.

1 13. A method for treating fungal infection in a mammal
2 in need of such treatment comprising administering to said mammal
3 an anti-fungal effective amount of a composition as defined in
4 claim 1.

1 14. A method for treating a tumor according to claim
2 11, wherein said tumor comprises a tumor of the prostate.

1 15. A method for treating a tumor according to claim
2 11, wherein said tumor comprises a tumor of the pancreas.

1 16. A method for treating a tumor according to claim
2 11, wherein said tumor comprises a tumor of the bladder.

1 17. A method for treating a fungal infection according
2 to claim 13, wherein said fungal infection comprises an infection
3 by *candida albicans*.

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US94/06653

A. CLASSIFICATION OF SUBJECT MATTER

IPC(5) : A61K 31/01, 31/02, 31/015, 31/19, 31/045, 31/265.

US CL : 514/574, 724, 763, 762, 512, 546, 547.

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

U.S. : 514/574, 724, 763, 762, 512, 546, 547.

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched
NONE

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

APS

SEARCH TERMS: ACONITIC, HEPATITIS, INFECTION, SORBIC, CROTONIC, MALIC.

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	US, A, 4,828,912 (HOSSAIN ET AL.) 09 MAY 1989, ENTIRE DOCUMENT.	1-17

☐ Further documents are listed in the continuation of Box C. ☐ See patent family annex.

Special categories of cited documents:		T	later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
A	document defining the general state of the art which is not considered to be of particular relevance		
E	earlier document published on or after the international filing date	*X*	document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
L	document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	*Y*	document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
O	document referring to an oral disclosure, use, exhibition or other means		
P	document published prior to the international filing date but later than the priority date claimed	*&*	document member of the same patent family

Date of the actual completion of the international search

25 AUGUST 1994

Date of mailing of the international search report

SEP 02 1994

Name and mailing address of the ISA/US
Commissioner of Patents and Trademarks
Box PCT
Washington, D.C. 20231

Facsimile No. (703) 305-3230

Authorized officer

THEODORE J. CRIARES

Telephone No. (703) 308-1235

THIS PAGE BLANK (USPTO)